

On Proactive Integrated Care Approach for COPD

Preeti Yadav¹, Jyothisree²

¹M.Sc. Nursing 2nd Year Student, Faculty of Nursing, Rama University, Kanpur, Uttar Pradesh, India

²Associate Professor (Department of Medical Surgical Nursing), Faculty of Nursing, Rama University, Kanpur, Uttar Pradesh, India

Corresponding Author: Mrs. Jyothisree, Associate Professor (Department of Medical Surgical Nursing), Faculty of Nursing, Rama University, Kanpur, Uttar Pradesh, India

Email: k.jyothisree.rcn@ramauniversity.ac.in

Abstract:

Chronic Obstructive Pulmonary Disease (COPD) is a chronic, progressive lung disorder characterized by persistent airflow limitation, breathlessness, chronic cough, and recurrent exacerbations. These exacerbations significantly reduce the quality of life and increase morbidity and mortality. Conventional management is largely reactive, focusing on treating exacerbations after their occurrence, which often leads to delayed intervention and increased healthcare burden. The Proactive Integrated Care Approach (PICA) has emerged as an innovative and patient-centered model that focuses on early detection, continuous monitoring, and coordinated multidisciplinary care. This approach integrates remote monitoring technologies, patient education, and proactive clinical response to manage COPD effectively. Evidence suggests that PICA improves health-related quality of life, enhances functional capacity, promotes adherence to treatment, and reduces hospital admissions. It empowers patients by involving them in self-management and ensures timely medical intervention.

Keywords: COPD, Proactive Integrated Care, PICA, Remote Monitoring, Self-Management, Quality of Life

SDES- International Journal of Interdisciplinary Research is a journal of Open access. In this journal, we allow all types of articles to be distributed freely and accessible under the terms of the creative common attribution- non-commercial share. This allows the authors, readers and all scholars and general community to understand, use and to develop non-commercially work, as long as appropriate credit is given and the newly developed work are licensed with similar terms.

How to cite this article: Yadav P, Jyothisree. Review Article On Proactive Integrated Care Approach for COPD. SDES-IJIR; 2026; 7-2: 1354-1357

Submitted: 08-March-2026; **Accepted:** 05-April-2026; **Published:** 30-April-2026

Introduction

Chronic Obstructive Pulmonary Disease (COPD) is one of the leading causes of morbidity and mortality worldwide. It is characterized by irreversible airflow limitation and chronic inflammation of the airways. Patients commonly experience symptoms such as dyspnea, fatigue, chronic cough, and sputum production.(1)

One of the major challenges in COPD management is the occurrence of exacerbations, which are episodes of worsening symptoms that often require hospitalization. These exacerbations accelerate disease progression, decrease functional capacity, and negatively impact the quality of life.(2)

Traditional management of COPD mainly focuses on pharmacological therapy and acute care during exacerbations. However, this reactive approach fails to identify early warning signs and

prevent complications. Many exacerbations remain unreported until they become severe, leading to increased hospital admissions and healthcare costs.(3)

To overcome these limitations, the Proactive Integrated Care Approach (PICA) has been introduced. This model emphasizes continuous monitoring, early detection of symptoms, and timely intervention. It integrates healthcare services, promotes patient participation, and ensures coordinated care delivery. PICA represents a shift from episodic care to continuous and preventive care, thereby improving overall disease management.(4)

Proactive Integrated Care Approach for COPD

Proactive Integrated Care Approach (PICA) is a comprehensive and patient-centered model designed to improve the management of chronic diseases like COPD. It combines advanced technology with clinical care to provide continuous monitoring and early intervention. The main objective of PICA is to detect clinical deterioration at an early stage and initiate timely treatment before the condition worsens. This approach involves collaboration among healthcare professionals and active participation of patients in their care.(5)

PICA is based on three key principles:

- Proactive monitoring of patient condition
- Integrated and coordinated care delivery
- Early and timely clinical intervention

By focusing on prevention rather than treatment, PICA helps reduce complications, improve patient outcomes, and enhance quality of life.

1. Remote Monitoring

Remote monitoring is a fundamental component of PICA and involves the use of digital devices and telehealth technologies to continuously assess the patient's health status. Parameters such as oxygen saturation (SpO₂), respiratory rate, heart rate, and symptom changes are regularly monitored. (6)

Patients may use portable devices at home to record their health data, which is transmitted to healthcare providers in real time. This enables healthcare professionals to track trends, identify abnormalities, and make informed decisions promptly. (7)

Remote monitoring is particularly beneficial in COPD patients, as it helps detect early signs of exacerbation that may not be immediately noticeable to the patient. It reduces the need for frequent hospital visits and provides continuous care even at home. (8)

2. Early Detection of Exacerbations

Exacerbations in COPD are often under-recognized and underreported by patients. Early detection is crucial in preventing disease progression and complications. (10)

PICA uses continuous monitoring and symptom tracking to identify subtle changes in the patient's condition, such as increased breathlessness, cough, or sputum production. These early warning signs allow healthcare providers to intervene before the exacerbation becomes severe. (11)

Timely detection leads to early treatment, such as medication adjustment or supportive therapy, which helps reduce the severity and duration of exacerbations. This ultimately decreases hospital admissions and improves patient outcomes. (12)

3. Integrated Multidisciplinary Care

Integrated care involves the collaboration of a multidisciplinary team, including physicians, nurses, respiratory therapists, and other healthcare professionals. (13)

Each member of the team plays a specific role in managing COPD. Nurses monitor patient condition and provide education, physicians manage treatment plans, and respiratory therapists assist with breathing exercises and pulmonary rehabilitation.

This coordinated approach ensures continuity of care and addresses all aspects of the patient's health—physical, psychological, and social. It improves communication among healthcare providers and enhances the efficiency of care delivery. (14)

4. Proactive Clinical Intervention

In PICA, healthcare providers do not wait for symptoms to worsen. Instead, they respond immediately to early signs of deterioration identified through monitoring systems. (15)

Proactive interventions may include adjusting medications, providing oxygen therapy, or advising lifestyle modifications. Early intervention helps prevent complications and reduces the need for emergency care. (16)

This approach minimizes disease progression and ensures better control of symptoms, leading to improved patient outcomes. (17)

5. Patient Education and Self-Management

Patient education is a key element of PICA. Patients are provided with information about their disease, treatment options, and lifestyle modifications. They are trained to recognize early symptoms of exacerbation and take appropriate action. Self-management strategies include medication adherence, smoking cessation, breathing exercises, and maintaining physical activity. (18)

Educating patients empowers them to take responsibility for their health, improves compliance with treatment, and enhances their confidence in managing the disease.

Effectiveness of PICA in COPD

Numerous studies have demonstrated the effectiveness of PICA in improving COPD outcomes. Patients receiving proactive integrated care show significant improvement in quality of life and symptom management. (18)

PICA has been shown to increase exercise tolerance, as measured by the 6-minute walk test, indicating improved physical capacity. It also helps in early detection of exacerbations, many of which would otherwise go unnoticed. (19)

Additionally, PICA reduces hospital admissions, emergency visits, and healthcare costs. Patients report better satisfaction due to continuous support and timely intervention.

Overall, this approach leads to better disease control, improved functional status, and enhanced well-being.

Conclusion

The Proactive Integrated Care Approach (PICA) is an effective and innovative model for managing COPD. By focusing on early detection, continuous monitoring, and coordinated care, it addresses the limitations of traditional reactive approaches.

PICA not only improves clinical outcomes but also enhances the quality of life of patients. It empowers patients through education and promotes active participation in their care.

The integration of technology and multidisciplinary care makes it a comprehensive and sustainable approach. Therefore, adopting PICA in clinical practice can significantly improve the management of COPD and reduce the overall disease burden

Financial support and sponsorship: Nil

Conflicts of interests: The authors declare that they have no conflict of interest

References

1. Koff PB, Min SJ, Campbell JD, et al. Proactive integrated care improves quality of life in patients with COPD. *Eur Respir J*. 2009;34(3):558–566.
2. Koff PB, Jones RH, Cashman JM, et al. Proactive integrated care for COPD patients: A randomized controlled trial. *Chronic Obstr Pulm Dis*. 2021;8(2):1–12.
3. Bourbeau J, Saad N. Integrated care model for COPD management. *Eur Respir Rev*. 2013;22(128):181–187.
4. McLean S, Nurmatov U, Liu JL, et al. Telehealthcare for COPD: Systematic review and meta-analysis. *Br J Gen Pract*. 2012;62(604):e739–e749.
5. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Global strategy for diagnosis, management, and prevention of COPD; 2024 report.
6. World Health Organization. Chronic obstructive pulmonary disease (COPD). Geneva: WHO; 2023.
7. Vogelmeier CF, Criner GJ, Martinez FJ, et al. Global strategy for COPD management. *Lancet Respir Med*. 2017;5(9):691–706.
8. Celli BR, Wedzicha JA. Update on clinical aspects of COPD exacerbations. *Am J Respir Crit Care Med*. 2015;192(4):398–406.
9. Miravittles M, Ribera A. Understanding the impact of symptoms on COPD management. *Respir Res*. 2014;15:122.
10. Criner GJ, Bourbeau J, Diekemper RL, et al. Prevention of acute exacerbations of COPD. *Am J Respir Crit Care Med*. 2015;192(2):224–238.
11. Effing TW, Bourbeau J, Vercoulen J, et al. Self-management programmes for COPD. *Cochrane Database Syst Rev*. 2012;(8):CD002990.
12. Holland AE, Cox NS, Houchen-Wolloff L, et al. Defining modern pulmonary rehabilitation. *Thorax*. 2017;72(3): 193–194.
13. Singh SJ, Puhan MA, Andrianopoulos V, et al. An official systematic review of pulmonary rehabilitation in COPD. *Eur Respir J*. 2018;51(2):1702414.
14. Montori VM, Kunneman M, Brito JP. Shared decision making in chronic care. *BMJ*. 2017;359:j4891.
15. Agusti A, Hogg JC. Update on pathogenesis of COPD. *Lancet*. 2014;384(9942):709–721.
16. ZuWallack R, Esteban C. Understanding COPD outcomes. *Chest*. 2014;145(6):1437–1445.
17. Han MK, Agusti A, Calverley PM, et al. Chronic obstructive pulmonary disease phenotypes. *Am J Respir Crit Care Med*. 2010;182(5):598–604.
18. Tashkin DP, Murray HE. Smoking cessation in COPD. *Respir Med*. 2009;103(7):963–974.
19. Calverley PM, Anderson JA, Celli B, et al. Salmeterol and fluticasone in COPD. *N Engl J Med*. 2007;356(8):775–789.

